

Vertical distribution of mesozooplankton and ichthyoplankton communities in the South-western Atlantic Ocean (23° 14' 1'' S 40° 42' 19'' W)

ANA C. T. BONECKER¹, CRISTINA DE O. DIAS¹, MARCIA S. DE CASTRO¹, PEDRO F. DE CARVALHO¹, ADRIANA V. ARAUJO¹, RODOLFO PARANHOS², ANDERSON S. CABRAL² AND SERGIO L. C. BONECKER¹

¹Universidade Federal do Rio de Janeiro, Instituto de Biologia, Departamento de Zoologia, Av. Carlos Chagas Filho, 373 – Prédio do CCS, Bloco A, Sala A0-084, Ilha do Fundão – 21.941-902, Rio de Janeiro, RJ, Brasil, ²Universidade Federal do Rio de Janeiro, Laboratório de Hidrobiologia, Departamento de Biologia Marinha, Instituto de Biologia, Av. Carlos Chagas Filho, 373 – Prédio do CCS, Bloco A, Cidade Universitária, Ilha do Fundão – 21.941-902, Rio de Janeiro, Brasil

*A study was conducted over eight consecutive days in February 2010 in which daily variations in the vertical distributions of heterotrophic bacteria, mesozooplankton and ichthyoplankton at 1–1200 m in the South-western Atlantic Ocean were investigated. Diurnal and nocturnal samples were collected at an oceanographic station at four regional depths: Tropical Water (TW) (1 m), South Atlantic Central Water (SACW) (250 m), Antarctic Intermediate Water (AAIW) (800 m) and Upper Circumpolar Deep Water (UCDW) (1200 m). Bacterial, mesozooplankton and larval fish densities significantly differed between sample depths but not between sampling tow times. In total, 154 zooplankton species and 18 larval fish species were identified. The highest number of taxa was obtained from the night-time TW trawls. This depth zone had the highest densities of mesozooplankton, larval fish and bacterioplankton (auto and heterotrophic), associated with the highest temperature and salinity and the lowest inorganic nutrient concentrations. Two sample groups were identified based on their mesozooplankton and larval fish compositions: night-time TW and other water masses (daytime TW, SACW, AAIW and UCDW). Thirty-two indicator species were detected in night-time TW. The copepod *Nullosetigera impar* was, to the best of our knowledge, identified for the first time on the Brazilian coast. Our results showed significant variability in the abundance and vertical distribution of mesozooplankton, bacterioplankton and larval fish along the water column in an oceanic area. We have provided new data and insights on the composition and vertical distribution of mesozooplankton, larval fish and bacterioplankton in deep waters in the South-western Atlantic Ocean.*

Keywords: mesozooplankton, fish larvae, bacterioplankton, vertical distribution, South-west Atlantic Ocean

Submitted 29 March 2017; accepted 22 November 2017

INTRODUCTION

Differences in vertical distribution are commonly observed in marine bacteria, mesozooplankton and ichthyoplankton (Tanaka & Rassoulzadegan, 2002; Yamaguchi *et al.*, 2002; Brugnano *et al.*, 2012; Siokou *et al.*, 2013; de Macedo-Soares *et al.*, 2014; Isla *et al.*, 2015; Munk *et al.*, 2015; Rodriguez *et al.*, 2015). The heterogeneous distribution of aquatic organisms has been studied since the early 20th century (Maycas *et al.*, 1999; Tiberti & Iacobuzio, 2013). Investigations have also been made concerning their diurnal movements in the water column (Roe, 1974; Krause & Radach, 1989). The vertical distribution of marine organisms is correlated with their physiological responses to several biological and physical factors. The former include visual predator avoidance, ontogenetic transport and dispersion regulation, and resource

competition (Cha *et al.*, 1994; Hill, 1998; Williamson *et al.*, 2011; Jung-Hoon *et al.*, 2013; Palomares-Garcia *et al.*, 2013; Harris *et al.*, 2014). Environmental factors include the hydrographic structure of the water column, light intensity, water temperature, salinity, density, dissolved oxygen (DO), current speed and current direction (Cha *et al.*, 1994; Williamson *et al.*, 2011; Brugnano *et al.*, 2012; Jung-Hoon *et al.*, 2013). Some mesozooplankton such as the Chaetognatha and Euphausiacea and larval fish families like Myctophidae and Gonostomatidae respond to light and other stimuli and are active migrators (Richards, 2006; Lie *et al.*, 2012; Sogawa *et al.*, 2016). The diel vertical migration of zooplankton is considered an anti-predatory response. It is triggered by changes in light intensity, but other factors are also involved in its regulation (Frost & Bollens, 1992; Hays, 2003; Pearre, 2003; Isla *et al.*, 2015). Zooplankton passively contribute to the carbon interchange along the water column with their own biomass when they die and sink along with faeces, mucous feeding webs, exoskeletons and carcasses (Angel, 2003; Conley & Hopkins, 2004; Castro *et al.*, 2010; Mayzaud & Pakhomov, 2014; Steinberg & Landry,

Corresponding author:
A.C.T. Bonecker
Email: ana@biologia.ufrj.br

2017). The vertical migration of zooplankton is an important component of the biological pump between the ocean surface and deeper waters (Steinberg & Landry, 2017).

The western boundary current system is located on the continental slope of Brazil between latitudes 22°S and 30°S. The upper part of this system is the Brazil Current flowing south-west towards the subtropical South Atlantic gyre (Peterson & Stramma, 1991). The lower part of the system consists of the Antarctic Intermediate Water (AAIW) with a variable circulation pattern along the Brazilian coast (Boebel *et al.*, 1997; Müller *et al.*, 1998). The circulation has intense mesoscale activity (Gabioux, 2008) with meandering, cyclonic and anticyclonic structures (Campos *et al.*, 1995, 1996).

The complexity of the water mass circulation has a profound impact on the natural resource diversity and ecological vulnerability of some marine areas (Gonzalez-Silvera *et al.*, 2004). Some studies reported on the vertical distribution of mesozooplankton and ichthyoplankton in the South-west Atlantic Ocean (Berasategui *et al.*, 2006; Dias *et al.*, 2010; Bonecker *et al.*, 2012, 2014a, b). None, however, addressed diurnal variations in vertical distribution. The aim of the present study was to examine the distribution and abundance of mesozooplankton and larval fish along the water column to a depth of 1200 m and to correlate the plankton with environmental variables. We expected mesozooplankton distribution to vary with day/night period and water mass. The results of this study help to elucidate the diel vertical migration patterns of mesozooplankton and larval fish in a tropical oceanic region.

METHODS

Study area

The northern region of Rio de Janeiro State has five water masses (Figure 1). The nutrient-poor Tropical Water (TW; temperature (T) >20 °C and salinity (S) >36.20) and the South Atlantic Central Water (SACW; 8.72 °C < T < 20 °C and 34.66 < S < 36.20) are found in the upper water column layers (Figure 1). At deeper levels, there are the cold waters of the Antarctic Intermediate Water (AAIW; 3.46 °C < T < 8.72 °C and 34.42 < S < 34.66), the Upper Circumpolar Deep Water (UCDW; 3.31 °C < T < 3.46 °C and 34.42 < S < 34.66) and the North Atlantic Deep Water (NADW; 2.04 °C < T < 3.31 °C and 34.59 < S < 34.87) (Mémery *et al.*, 2000; Silveira, 2007; Bonecker *et al.*, 2012, 2014b).

Sampling collection and processing

Water, heterotrophic bacteria and zooplankton samples were collected at an oceanographic station (23°14'1"S 40°42'19"W) at four depths corresponding to the previously defined water masses: TW (1 m), SACW (250 m), AAIW (800 m) and UCDW (1200 m) (Figure 2). These sampling depths represent each water mass nucleus. Samples were collected for eight consecutive days in the rainy season (February 2010) in the daytime (06:57 to 16:15 h) and at night (19:45 to 05:32 h) (GMT + 3). Fourteen samples were collected at night and 18 during the day. Diurnal samples were considered replicates, and the same was done with the samples collected at night.

Water temperature and salinity were determined with a Rosette system fitted with a CTD profiler (Sea-Bird




Fig. 1. Salinity and temperature of the five water masses (0–3260 m) in the Campos Basin, central Brazilian coast. Modified from Bonecker *et al.* (2014b). Solid line, temperature; dashed line, salinity. SS, subsurface water; SACW, South Atlantic Central Water; AAIW, Antarctic Intermediate Water; UCDW, Upper Circumpolar Deep Water; NADW, North Atlantic Deep Water.

Electronics, Inc., Bellevue, WA, USA). Water samples were collected using a GO-FLO bottle (General Oceanics, Miami, FL, USA) for the analysis of inorganic nutrients (nitrate, silicate and orthophosphate). These were determined using standard oceanographic methods (Grasshoff *et al.*, 1999). DO in the water column was measured continuously using a sensor coupling in the CTD. Temperature, salinity and samples for nutrient analysis and DO were obtained at each collection. A total of 32 data points was determined for each variable. Detailed methodologies and discussions on the hydrochemistry of the study area have been presented elsewhere (Rodrigues *et al.*, 2014; Bonecker *et al.*, 2014b; Dias *et al.*, 2015; Suzuki *et al.*, 2015).

Samples for the assessment of the abundance of bacteria (autotrophic and heterotrophic) were collected in Niskin bottles and then transferred to 2-ml Eppendorf vials. They were fixed *in situ* with a mixture of 1% v/v paraformaldehyde and 0.05% v/v glutaraldehyde, frozen in liquid nitrogen, and maintained there until analysis (Gasol & del Giorgio, 2000; Andrade *et al.*, 2003). In the laboratory, aliquots of heterotrophic bacterial samples were stained with SYBR Green I (Molecular Probes, Eugene, OR, USA) at a final concentration of 5×10^{-5} of the commercial stock solution (Gasol & del Giorgio, 2000; Andrade *et al.*, 2003). They were analysed using a FACSCalibur flow cytometer (BD Biosciences, San Jose, CA, USA) equipped with a 488-nm argon laser. Prokaryotic heterotrophic cells with high- or low nucleic




Fig. 2. Sampling station off the central Brazilian coast surveyed in this study. Lines indicate isobaths.

acid content (HNA and LNA, respectively) were detected, identified and quantified based on their signatures in a side scatter plot (X -axis; correlated by size) against green fluorescence (Y -axis; SYBR Green I staining; correlated by nucleic acid content) (Gasol & del Giorgio, 2000; Andrade *et al.*, 2003). Autotrophic bacteria (*Synechococcus* and *Phrochlorococcus*) were analysed with the same instrument using the autofluorescence of pigments (chlorophyll on the red detector, and phycoerythrin on the orange detector). The abundance of heterotrophic bacteria was calculated by subtracting the autotrophic contribution (Gasol & del Giorgio, 2000).

Horizontal hauls were performed using a Midi-type Hydro-Bios MultiNet® (aperture 0.25 m^2) fitted with a set of two nets (mesh apertures 200 and $500 \mu\text{m}$) used to sample each water mass separately to prevent cross-contamination. At each predetermined depth, hauls were performed at a speed of two knots with an open–close mechanism operated by electronically transmitted commands. The MultiNet was equipped with a depth sensor. The haul depth was controlled during the entire procedure to ensure that the net was towed horizontally. In TW and SACW, hauls were run for 10 min whereas in AAIW and UCDW the nets were towed for 15 min due to the low organism density in the deeper waters. Water volume and haul depth data were transmitted in real time to a computer on board the ship. Filtration efficiency and water volume were measured using flowmeters. The average water volumes filtered through the

$200\text{-}\mu\text{m}$ mesh were $116.0 \pm 40.4 \text{ m}^{-3}$, $89.0 \pm 31.3 \text{ m}^{-3}$, $126.8 \pm 42.1 \text{ m}^{-3}$ and $104.5 \pm 28.8 \text{ m}^{-3}$ in TW, SACW, AAIW and UCDW, respectively. The mean water volumes filtered through the $500\text{-}\mu\text{m}$ net were $132.9 \pm 39.2 \text{ m}^{-3}$, $96.8 \pm 27.9 \text{ m}^{-3}$, $121.6 \pm 29.6 \text{ m}^{-3}$ and $112.0 \pm 30.2 \text{ m}^{-3}$ in TW, SACW, AAIW and UCDW, respectively.

The samples were immediately fixed in 4% v/v buffered formalin. Mesozooplankton were analysed in 29 samples collected with a $200\text{-}\mu\text{m}$ mesh net. A $500\text{-}\mu\text{m}$ net was used to harvest 32 larval fish samples. The difference in the numbers of mesozooplankton and larval fish samples was the result of logistical problems during the hauls. In the laboratory, the mesozooplankton were divided into subsamples using a Folsom's Plankton Sample Splitter (Hydro-Bios, Am Jägersberg, Altenholz, Germany) (McEwen *et al.*, 1954). The degree of subsampling was adjusted according to the organism density so that 100 individuals per taxonomic group were allocated to each sample. The lots were sorted and counted from fractions of $1/16$ – $1/1024$ in TW and from $1/1$ – $1/32$ in SACW. Organisms present in all AAIW and UCDW samples were completely sorted. The samples collected using the $500\text{-}\mu\text{m}$ mesh were entirely sorted for larval fish. The mesozooplankton and larval fish catches were standardized to the number of individuals m^{-3} . The mesozooplankton groups (Mollusca: Cephalopoda, Branchiopoda, Copepoda, Euphausiacea, Decapoda, Chaetognatha, Appendicularia, Salpida and Doliolida) and the larval fish were identified to the lowest taxonomic level possible using

specialized literature (Boltovskoy, 1999; Bonecker, 2006; Bonecker & Castro, 2006; Richards, 2006; Bonecker *et al.*, 2014a).

Data analysis

Daytime/night-time abundance, vertical density differences, and the interactions between these two factors were tested using generalized linear models (GLM) with gamma family (dispersion = 1). We used the densities of total mesozooplankton, mesozooplankton groups with >5% relative abundance (Copepoda, Branchiopoda, Mollusca, Euphausiacea, Chaetognatha, Appendicularia, Doliolida, Salpida), total larval fish, larval fish families comprising >10% of the total catch, heterotrophic bacteria, %HNA bacteria and %LNA bacteria. Because autotrophic bacterioplankton were only detected in the surface layers, they were not included in the data analyses. A low additive constant of one was added to the density data to eliminate zero values in the matrix because the gamma family does not allow for them. The results were considered significant only when $P < 0.05$. The analyses were performed using R v.2.12.1 (R Development Core Team, 2010; <http://www.r-project.org>).

Principal component analysis (PCA) was used to define the similarities between the samples according to environmental descriptors (continuous variables) and to define how the observed patterns relate to the environment. The environmental descriptors were temperature, salinity, DO, nitrate, silicate and orthophosphate. The environmental variables were standardized and normalized for the different water masses (TW, SACW, AAIW and UCDW) before the PCA was run. Mesozooplankton, larval fish, heterotrophic bacterial abundance, %HNA and %LNA were added as categorical supplements. The correlation matrix was used to calculate eigenvectors and principal components (PC) which were ranked in the order of significance. The broken-stick method was used as a stopping rule in the PCA (Jackson, 1993). This analysis was performed using PCORD v.5 (McCune & Mefford, 1999). Scores of the retained PCA axes were used as new variables to determine whether the environmental data varied with depth by using Gaussian ANOVA. The results were considered significant only when $P < 0.05$. The analyses were performed using R v.2.12.1 (R Development Core Team, 2010; <http://www.r-project.org>).

A hierarchical agglomerative cluster analysis was computed with the Sorenson similarity index based on the presence or absence of mesozooplankton groups with >5% relative abundance. The matrix was based on 90 species and 29 samples. A dendrogram was constructed using the weighted pair group of arithmetic averages method. Similarity percentages (SIMPER;

Clarke & Warwick, 1994) and the Euclidean distance index were used to identify the species that contributed the most to the average similarity and dissimilarity within each group. All analyses were run with PRIMER v.6.0 according to the method described by Clarke & Warwick (1994).

For the PCA and cluster analyses, three samples were excluded because they did not match the others obtained from the mesozooplankton collection using the 200- μm net deployment.

We performed an indicator species analysis (ISA; Dufrene & Legendre, 1997) to include species abundance data and identify the indicator taxa for each water mass (TW, SACW, AAIW and UCDW) and time period (day, night). The indicator value (IndVal) of a taxon is the product of the relative frequency of its occurrence and its relative average abundance in previously defined groups multiplied by 100. A 100% ISA index value is obtained when all representatives of a species are found within a single sample group and the species occurs in all the samples of that group. A species was considered an indicator of a particular water mass when its IndVal was >70% and significantly higher than that compared to one thousand random samples of plots with the same number of species occurrences. Species for which the IndVal was <70% were considered detectors (Van Rensburg *et al.*, 1999; McGeoch *et al.*, 2002). These values were statistically analysed using the Monte Carlo test to establish reliable significance levels ($P < 0.05$). The analysis was performed using PCORD v.5.

RESULTS

Environmental data

Temperature, salinity and DO were distributed along a typical oceanic gradient towards deeper waters, being more variable at the surface (Table 1). The mean water temperature ranged from 3.4 °C (UCDW) to 28.2 °C (TW). Salinity was relatively stable in deeper waters (mean < 34.6), as did DO (mean < 4.5 mg l⁻¹). Nutrient concentrations were highest in deeper waters (mean > 30 $\mu\text{mol l}^{-1}$ for nitrate, > 17 $\mu\text{mol l}^{-1}$ for silicate, and > 1.6 $\mu\text{mol l}^{-1}$ for orthophosphate (Table 1).

Bacteria, mesozooplankton and larval fish distributions

The autotrophic bacterioplankton group was dominated by *Prochlorococcus* (between 2.7 and 8.4 $\times 10^4$ cells ml⁻¹), with

Table 1. Means and standard deviations of the environmental variables (temperature, °C; salinity, dissolved oxygen (DO), ml l⁻¹; nitrate, silicate, and orthophosphate, $\mu\text{mol l}^{-1}$) measured in TW, SACW, AAIW and UCDW in the daytime and at night. N, number of samples.

Water mass/period	Temperature	Salinity	DO	Nitrate	Silicate	Orthophosphate	N
TW night	28.20 \pm 0.08	36.96 \pm 0.03	4.24 \pm 0.06	1.75 \pm 0.54	0.78 \pm 0.29	0.03 \pm 0.01	4
TW day	28.19 \pm 0.09	36.98 \pm 0.05	4.17 \pm 0.10	1.82 \pm 0.35	0.99 \pm 0.03	0.03 \pm 0.01	4
SACW night	14.86 \pm 0.12	35.46 \pm 0.02	4.47 \pm 0.03	4.17 \pm 0.81	2.08 \pm 0.15	0.45 \pm 0.05	3
SACW day	14.85 \pm 0.12	35.45 \pm 0.02	4.48 \pm 0.01	3.06 \pm 1.00	2.20 \pm 0.59	0.48 \pm 0.05	5
AAIW night	4.75 \pm 0.19	34.35 \pm 0.03	4.42 \pm 0.18	30.64 \pm 10.34	17.43 \pm 4.71	1.67 \pm 0.24	4
AAIW day	4.83 \pm 0.23	34.36 \pm 0.02	4.36 \pm 0.14	33.46 \pm 9.75	19.17 \pm 4.47	1.64 \pm 0.22	4
UCDW night	3.42 \pm 0.06	34.53 \pm 0.01	4.07 \pm 0.01	37.07 \pm 2.67	35.71 \pm 3.06	1.90 \pm 0.03	3
UCDW day	3.43 \pm 0.10	34.53 \pm 0.01	4.06 \pm 0.01	32.47 \pm 7.11	38.31 \pm 6.56	1.95 \pm 0.19	5

smaller contribution of *Synechococcus* (between 0.8 and 3.5×10^3 cells ml^{-1}). The contribution of autotrophic bacteria to microbial biomass in the surface waters (TW) ranged from 7 to 18%, whereas its contribution was negligible below 200 m. The abundance of heterotrophic bacteria decreased one order of magnitude from the surface (1.5 to 5.7×10^5 cells l^{-1}) to deep waters (3.0 to 6.0×10^4 cells ml^{-1} ; Figure 3), and these differences were significant ($P < 0.05$). No interaction was observed between the water masses and day/night period. Among heterotrophic bacterial subgroups, LNA bacteria dominated the euphotic zone constituting 80–90% of the total counts. Total heterotrophic bacteria decreased towards deep waters, however an increase in % HNA cells were observed and such bigger cells dominated microbial biomass at deep waters (Table 2). Similar to the results of heterotrophic bacterial abundance, significant differences were observed only for water masses ($P < 0.05$). The distributions of mesozooplankton and larval fish within each water mass followed the same pattern as that observed for bacteria. The highest abundance of total mesozooplankton and larval fish was observed in TW and SACW (Figure 4). The densities of each mesozooplankton group were highest in TW at night except for the Branchiopoda, which were more numerous in the daytime (Table 3). Copepoda and Chaetognatha were present in all water masses and in both sampling periods whereas Branchiopoda and Salpida occurred only in TW and SACW (Table 3). Gonostomatidae (TW to UCDW) and Myctophidae (TW to AAIW) larval fish showed wide vertical distributions whereas scombrids were detected only in TW. Myctophidae and Scombridae densities were highest in TW during the night whereas gonostomatids occurred mainly in AAIW at night (Table 3). Copepoda was the most abundant group (72–95%), followed by mollusc larvae (1–18%). The Gonostomatidae predominated (3–100%), and they were the only larval fish representatives in UCDW (Table 3).

Despite what was described earlier no interaction was observed between the water masses and day/night period in terms of total mesozooplankton and larval fish abundance. Significant differences were detected only among the water masses ($P < 0.05$). Nevertheless, individual analyses of the most abundant mesozooplankton and larval fish families

revealed significant ($P < 0.05$) differences in the interactions between vertical distribution and daytime/night-time abundance, but only for the Euphausiacea, Chaetognatha, Salpida and Doliolida.

A total of 154 mesozooplankton species (two molluscs/cephalopods, two branchiopods, 112 copepods, three larval decapods, 12 euphausiids, 10 chaetognaths, eight appendicularians, two doliolids, and three salpids) and 18 larval fish species were identified. The highest number of taxa was obtained in TW at night (Figure 5). To the best of our knowledge, the present study is the first to report on the copepod *Nullosetigera impar* in the South-west Atlantic Ocean. This species occurred in AAIW during both time periods. Molluscs were the second most numerous group in TW (Table 3).

Influence of environmental variables

The first two axes of the PCA performed on the environmental factors accounted for 96% of the total variance. Only PC 1 (eigenvalue = 2.45) was retained in the analyses to explain the data variability (76%). The PCA showed that the four water masses were separated where the samples were drawn (axis 1). Temperature and salinity accounted for positive separation (0.44 and 0.43, respectively), whereas orthophosphate, nitrate and silicate explained negative separation (−0.47, −0.44 and −0.43, respectively). TW and SACW (right side of the plot) were influenced by the highest temperatures and salinities and by the lowest orthophosphate, nitrate and silicate concentrations. The deeper water masses (AAIW and UCDW; left side of the plot) showed the opposite trend to the shallower water masses (Figure 6). The scores of axis 1 indicated significant differences between depths (GLM; $F = 19.18$; $df = 3$; $P < 0.05$). Therefore, the variables related to axis 1 varied depending on the water mass characteristics. In TW, the mesozooplankton groups, larval fish, heterotrophic bacterial abundance and LNA bacteria increased as temperature and salinity increased and as inorganic nutrient (nitrate, silicate, and orthophosphate) concentrations decreased. Conversely, HNA bacteria increased with inorganic nutrient concentration in AAIW and UCDW.




Fig. 3. Heterotrophic bacterial abundances collected in the daytime (white) and at night-time (black) in TW, SACW, AAIW and UCDW. Abundance is expressed as \log_{10} cells ml^{-1} . Consecutive sampling days are labelled from 1–8.

Table 2. Means and standard deviations of the heterotrophic bacteria groups (%HNA and %LNA) measured in TW, SACW, AAIW and UCDW in the daytime and at night.

Water mass/period	%HNA	%LNA
TW night	30 ± 22	70 ± 22
TW day	29 ± 15	71 ± 15
SACW night	39 ± 2	61 ± 2
SACW day	39 ± 3	61 ± 3
AAIW night	58 ± 17	42 ± 17
AAIW day	69 ± 10	31 ± 10
UCDW night	64 ± 27	36 ± 27
UCDW day	64 ± 10	36 ± 10

Mesozooplankton and larval fish communities

The cluster analysis showed two sample groups based on the composition of the mesozooplankton and larval fish

communities at an 80% similarity level. Group I consisted of the night-time TW samples and Group II comprised all other samples (Figure 7). The mesozooplankton and larval fish species contributing to group similarity are shown in Table 4 (SIMPER test). The night-time TW group was composed of 22 species of which 13 contributed 3.61% each: four larval fish, 11 copepods, one chaetognath, two appendicularians, one salp, one mollusc, one decapod and one euphausiid. Two larval fish, one mollusc, four copepods, one decapod larva and one euphausiid each contributed 4.82% to the formation of this group (Table 4).

The samples collected in the other water masses (daytime TW, SACW, AAIW and UCDW) included 76 species (Table 4). Among them, copepods were the most representative, with 51 species. There were 15 species (including nine copepods, one euphausiid, two appendicularians and three larval fish) in common between the night-time TW and the other water masses (Table 4).




Fig. 4. Total abundance of organisms collected during the day (white) and at night (black) in TW, SACW, AAIW and UCDW. Abundance is expressed as \log_{10} of the number of specimens m^{-3} for mesozooplankton (a) and larval fish (b). Consecutive sampling days are labelled from 1–8.

Table 3. Mean abundance, standard deviation (number of specimens m^{-3}), and relative abundance (%) of the most abundant mesozooplanktonic groups and larval fish families collected during the day and at night from TW, SACW, AAIW and UCDW. –, absence of the organisms listed.

Taxon	TW		SACW		AAIW		UCDW	
	Daytime	Night-time	Daytime	Night-time	Daytime	Night-time	Daytime	Night-time
Mesozooplankton								
Mollusca	1.76 ± 2.48 0.91%	161.16 ± 198.14 18.36%	2.02 ± 2.31 8.39%	0.11 ± 0.06 0.73%	–	–	0.01 ± 0.01 1.38%	–
Branchiopoda	1.11 ± 1.92 0.57%	0.05 ± 0.04 0.01%	0.003 ± 0.01 0.01%	0.04 ± 0.06 0.27%	–	–	–	–
Copepoda	181.37 ± 258.91 93.61%	635.36 ± 543.78 72.39%	17.76 ± 14.97 73.77%	14.27 ± 16.51 95.18%	0.70 ± 0.73 92.11%	0.82 ± 0.39 86.86%	0.67 ± 0.32 92.67%	0.63 ± 0.20 90.00%
Euphausiacea	0.02 ± 0.01 0.01%	5.33 ± 4.72 0.61%	0.03 ± 0.01 0.12%	0.03 ± 0.05 0.20%	–	–	0.003 ± 0.01 0.41%	–
Chaetognatha	0.59 ± 0.56 0.30%	16.81 ± 6.72 1.92%	0.51 ± 0.48 2.12%	0.39 ± 0.45 2.60%	0.01 ± 0.02 1.32%	0.03 ± 0.02 3.18%	0.01 ± 0.01 1.38%	0.02 ± 0.02 2.86%
Appendicularia	2.58 ± 3.02 1.33%	41.96 ± 16.92 4.78%	0.05 ± 0.06 0.21%	0.04 ± 0.02 0.27%	–	0.01 ± 0.02	0.01 ± 0.02 1.38%	–
Doliolida	0.13 ± 0.10 0.07%	1.28 ± 1.10 0.15%	0.002 ± 0.004 0.01%	0.003 ± 0.01 0.02%	–	0.004 ± 0.01 0.42%	–	–
Salpida	0.06 ± 0.08 0.03%	0.29 ± 0.25 0.03%	0.02 ± 0.03 0.08%	–	–	–	–	–
Other groups	6.13 ± 4.83 3.16%	15.45 ± 13.50 1.76%	3.68 ± 3.73 15.29%	0.11 ± 0.07 0.73%	0.05 ± 0.04 6.58%	0.08 ± 0.02 8.47%	0.02 ± 0.03 2.77%	0.05 ± 0.03 7.14%
Larval fish								
Gonostomatidae	0.26 ± 0.44 3.63%	0.92 ± 0.81 3.07%	–	–	1.06 ± 0.34 56.38%	1.99 ± 1.28 81.89%	0.58 ± 0.58 100.00%	–
Myctophidae	1.21 ± 0.95 16.88%	20.96 ± 8.37 69.96%	0.36 ± 0.52 50.00%	0.41 ± 0.58 18.14%	0.49 ± 0.99 20.06%	0.44 ± 0.88 18.11%	–	–
Scombridae	1.13 ± 1.95 15.76%	5.94 ± 3.65 19.83%	–	–	–	–	–	–
Other larval fish	4.57 ± 3.42 63.74%	2.14 ± 0.94 7.14%	0.36 ± 0.52 50.00%	1.85 ± 2.62 81.86%	0.33 ± 0.66 17.55%	–	–	–




Fig. 5. Number of mesozooplankton taxa collected during the daytime (white) and at night (black) from TW, SACW, AAIW and UCDW.

The ISA showed 32 indicator species exclusive to the night-time TW (Table 5). Two detector species were recorded in the daytime TW, and one each in the night-time TW, SACW, AAIW and UCDW (Table 5).

DISCUSSION

Oceanographic conditions

The hydrological variables measured in this study resembled those of previous experiments along the Brazilian coast and were typical of the oligotrophic oceanic region (Rezende *et al.*, 2007; Rodrigues *et al.*, 2014). The environmental variables described for the water column reflected the unique hydrological signatures of the water masses. The regional temperature and salinity data are characteristics of the water masses there (Niencheski *et al.*, 1999; Rezende *et al.*, 2007). These variables decreased from the subsurface (TW; depth 1 m) to the deep waters (UCDW; depth 1200 m). Average DO did not vary greatly in the study area and resembled other DO values obtained for this oceanic region of Brazil (Rezende *et al.*, 2007; Suzuki *et al.*, 2015). Inorganic nutrient




Fig. 7. Cluster analysis based on species composition in the samples collected during the daytime and at night-time from TW, SACW, AAIW and UCDW. The Sorensen-Dice coefficient and the average linkage method were used. Different groups indicate faunistic zones defined at 80% similarity. Data labels: N, night-time; D, daytime; TW, Tropical Water; SACW, South Atlantic Central Water; AAIW, Antarctic Intermediate Water; UCDW, Upper Circumpolar Deep Water.

concentrations showed a typical oceanic vertical distribution pattern: they increased from the surface to deeper waters. In the study area, nutrients are usually depleted in the surface waters (Rezende *et al.*, 2007; Rodrigues *et al.*, 2014; Suzuki *et al.*, 2015). This pattern is attributed to the high consumption of nutrients by the primary producers during photosynthesis in TW (Rodrigues *et al.*, 2014). The low nitrate, silicate and orthophosphate levels in TW are characteristic of the nutrient-poor oceanic waters carried by the Brazil Current (Rezende *et al.*, 2007; Alves *et al.*, 2014; Rodrigues *et al.*, 2014).




Fig. 6. PCA output used to summarize environmental and biological variables. Abiotic variables were as follows: temperature (Tem), salinity (Sal), dissolved oxygen (DO), nitrate (Nit), silicate (Sil) and orthophosphate (P-Inor). Mesozooplankton (Zoo), larval fish (LF), heterotrophic bacteria (Bac), LNA bacteria (LNA) and HNA bacteria (HNA) were added as categorical supplements. Samples collected from TW (black square, night; open square, day), SACW (black polygon, night; open polygon, day), AAIW (black circle, night; open circle, day), and UCDW (black triangle, night; open triangle, day) were arranged according to the first two principal components.

Table 4. Mesozooplankton species and their contribution (%) to the similarity of the communities determined by SIMPER analysis of the two groups formed by the cluster analysis. The group formed by SACW, AAIW and UCDW included samples collected during the day and at night.

TW night-time			TW daytime/SACW/AAIW/UCDW		
Average squared distance = 6.92%			Average squared distance = 10.37%		
Code	Species	(%)	Code	Species	(%)
L	<i>Coryphaena equiselis</i>	3.61	L	<i>Coryphaena equiselis</i>	0.39
L	<i>Lepidophanes guentheri</i>	3.61	Co	<i>Candacia pachydactyla</i>	0.39
Co	<i>Calocalanus pavoninus</i>	3.61	Co	<i>Centropages</i> sp.	0.39
Co	<i>Candacia pachydactyla</i>	3.61	Co	<i>Haloptilus acutifrons</i>	0.39
Co	<i>Centropages</i> sp.	3.61	Co	<i>Pleuromamma abdominalis</i>	0.39
Co	<i>Oithona similis</i>	3.61	Co	<i>Sapphirina nigromaculata</i>	0.39
Co	<i>Scolecithrix danae</i>	3.61	E	<i>Stylocheiron carinatum</i>	0.39
Co	<i>Spinocalanus</i> sp.	3.61	E	<i>Thysanopoda aequalis</i>	0.39
Co	<i>Sapphirina nigromaculata</i>	3.61	E	<i>Thysanopoda tricuspidata</i>	0.39
C	<i>Serratosagitta serratodentata</i>	3.61	C	<i>Kronitta pacifica</i>	0.39
A	<i>Fritillaria formica</i>	3.61	A	<i>Oikopleura cornutogastra</i>	0.39
A	<i>Oikopleura cornutogastra</i>	3.61	D	<i>Doliolletta gegenbauri</i>	0.39
S	<i>Brooksia rostrata</i>	3.61	L	<i>Dactylopterus volitans</i>	0.74
L	<i>Cyclothone braueri</i>	4.82	Co	<i>Aegisthus mucronatus</i>	0.74
L	<i>Dactylopterus volitans</i>	4.82	Co	<i>Aetideus giesbrechti</i>	0.74
M	<i>Abralia</i> sp.	4.82	Co	<i>Agetus typicus</i>	0.74
Co	<i>Lucicutia flavicornis</i>	4.82	Co	<i>Clausocalanus arcuicornis</i>	0.74
Co	<i>Onchocorycaeus giesbrechti</i>	4.82	Co	<i>Haloptilus austini</i>	0.74
Co	<i>Pleuromamma abdominalis</i>	4.82	Co	<i>Lucicutia ovalis</i>	0.74
Co	<i>Triconia</i> cf. <i>conifera</i>	4.82	Co	<i>Miracia efferata</i>	0.74
De	<i>Palaemon</i> sp.	4.82	Co	<i>Nullosetigera impar</i>	0.74
E	<i>Stylocheiron carinatum</i>	4.82	Co	<i>Oithona plumifera</i>	0.74

TW daytime/SACW/AAIW/UCDW					
Average squared distance = 10.37%					
Code	Species	(%)	Code	Species	(%)
Co	<i>Onchocorycaeus giesbrechti</i>	0.74	Co	<i>Lucicutia gaussae</i>	1.35
Co	<i>Scaphocalanus echinatus</i>	0.74	Co	<i>Paracalanus quasimodo</i>	1.35
Co	<i>Scaphocalanus elongatus</i>	0.74	Co	<i>Scolecithrix danae</i>	1.35
E	<i>Nematoscelis tenella</i>	0.74	A	<i>Oikopleura rufescens</i>	1.35
E	<i>Stylocheiron longicorne</i>	0.74	Co	<i>Lucicutia flavicornis</i>	1.61
C	<i>Pseudosagitta lyra</i>	0.74	Co	<i>Pareucalanus sewelli</i>	1.61
A	<i>Fritillaria formica</i>	0.74	Co	<i>Temoropia mayumbaensis</i>	1.61
S	<i>Salpa fusiformis</i>	0.74	C	<i>Decipisagitta sibogae</i>	1.61
S	<i>Thalia democratica</i>	0.74	A	<i>Oikopleura fusiformis</i>	1.61
Co	<i>Lophothrix</i> sp.	1.06	L	<i>Cyclothone braueri</i>	1.83
Co	<i>Lophothrix frontalis</i>	1.06	Co	<i>Acrocalanus longicornis</i>	1.83
Co	<i>Macrosetella gracilis</i>	1.06	Co	<i>Corycaeus speciosus</i>	1.83
Co	<i>Microsetella rosea</i>	1.06	Co	<i>Euaugaptilus</i> sp.	1.83
Co	<i>Nannocalanus minor</i>	1.06	Co	<i>Microsetella norvegica</i>	1.83
Co	<i>Oncaea venusta</i>	1.06	Co	<i>Oithona similis</i>	1.83
Co	<i>Paraheterorhabdus vipera</i>	1.06	Co	<i>Oncaea media</i>	1.83
Co	<i>Scaphocalanus curtus</i>	1.06	Co	<i>Oithona setigera</i>	2.02
Co	<i>Temora turbinata</i>	1.06	Co	<i>Undinula vulgaris</i>	2.02
Co	<i>Urocorycaeus lautus</i>	1.06	C	<i>Flaccisagitta enflata</i>	2.02
De	<i>Gennadas</i> sp.	1.06	D	<i>Doliolum nationalis</i>	2.02
De	<i>Lucifer typus</i>	1.06	Co	<i>Calanoides</i> cf. <i>carinatus</i>	2.31
C	<i>Caecosagitta macrocephala</i>	1.06	Co	<i>Rhincalanus cornutus</i>	2.31
B	<i>Pseudevadne tergestina</i>	1.35	Co	<i>Spinocalanus</i> sp.	2.31
Co	<i>Calocalanus pavo</i>	1.35	Co	<i>Temora styliifera</i>	2.31
Co	<i>Chirundina</i> sp.	1.35	Co	<i>Clausocalanus furcatus</i>	2.41
Co	<i>Clausocalanus brevipes</i>	1.35	Co	<i>Conaea rapax</i>	2.41
Co	<i>Clausocalanus mastigophorus</i>	1.35	C	<i>Parasagitta friderici</i>	2.41

B, Branchiopoda; Co, Copepoda; E, Euphausiacea; De, Decapoda; C, Chaetognatha; A, Appendicularia; S, Salpida; D, Doliolida; M, Mollusca; L, Larval fish.

Table 5. Indicator species, taxonomic group, water mass, period, IndVal (%) and Monte Carlo test significance (*P*).

Species	Group	Water mass/ period	IndVal	<i>P</i>
<i>Nannocalanus minor</i>	Copepoda	TW/night	100.0	0.0002
<i>Pterosagitta draco</i>	Chaetognatha	TW/night	100.0	0.0004
<i>Paracalanus quasimodo</i>	Copepoda	TW/night	100.0	0.0004
<i>Euphausia americana</i>	Euphausiacea	TW/night	100.0	0.0004
<i>Euphausia similis</i>	Euphausiacea	TW/night	100.0	0.0004
<i>Nematoscelis atlantica</i>	Euphausiacea	TW/night	100.0	0.0004
<i>Lepidophanes gausi</i>	Larval fish	TW/night	100.0	0.0004
<i>Clausocalanus furcatus</i>	Copepoda	TW/night	99.8	0.0006
<i>Temora stylifera</i>	Copepoda	TW/night	99.5	0.0018
<i>Krohnittia mutabii</i>	Chaetognatha	TW/night	99.2	0.0004
<i>Corycaeus speciosus</i>	Copepoda	TW/night	99.1	0.0004
<i>Oncaea venusta</i>	Copepoda	TW/night	98.8	0.0044
<i>Oikopleura rufescens</i>	Appendicularia	TW/night	98.5	0.0004
<i>Thysanopoda aequalis</i>	Euphausiacea	TW/night	98.1	0.0006
<i>Oncaea media</i>	Copepoda	TW/night	97.8	0.0046
<i>Acrocalanus longicornis</i>	Copepoda	TW/night	97.7	0.0118
<i>Farranula gracilis</i>	Copepoda	TW/night	97.5	0.0180
<i>Macrosetella gracilis</i>	Copepoda	TW/night	97.2	0.0338
<i>Lucifer typus</i>	Decapoda	TW/night	97.1	0.0004
<i>Undinula vulgaris</i>	Copepoda	TW/night	96.5	0.0072
<i>Flaccisagitta enflata</i>	Chaetognatha	TW/night	96.3	0.0004
<i>Oikopleura fusiformis</i>	Appendicularia	TW/night	94.2	0.0004
<i>Oikopleura longicauda</i>	Appendicularia	TW/night	93.3	0.0004
<i>Doliolum nationalis</i>	Thaliacea	TW/night	90.2	0.0004
<i>Parasagitta friderici</i>	Chaetognatha	TW/night	89.9	0.0002
<i>Salpa fusiformis</i>	Thaliacea	TW/night	89.8	0.0004
<i>Serratosagitta serratodentata</i>	Chaetognatha	TW/night	75.0	0.0064
<i>Brooksia rostrata</i>	Thaliacea	TW/night	75.0	0.0064
<i>Lepidophanes guentheri</i>	Larval fish	TW/night	75.0	0.0066
<i>Scolecithrix danae</i>	Copepoda	TW/night	74.7	0.0370
<i>Calocalanus pavo</i>	Copepoda	TW/night	73.8	0.0188
<i>Fritillaria formica</i>	Appendicularia	TW/night	72.6	0.0090
<i>Gennada</i> sp.	Decapoda	TW/day	66.9	0.0016
<i>Nematoscelis tenella</i>	Euphausiacea	SACW/ night	66.7	0.0234
<i>Cyclothone braueri</i>	Larval fish	AAIW/night	64.2	0.0090
<i>Dactylopterus volitans</i>	Larval fish	TW/Day	60.4	0.0250
<i>Thalia democratica</i>	Thaliacea	TW/night	55.8	0.0192
<i>Conaea rapax</i>	Copepoda	UDCW/ night	54.6	0.0090

Diurnal and vertical variations over an eight-day period

Previous microbiological studies in the South-west Atlantic Ocean were conducted mainly on surface water sampled during transatlantic cruises (Zubkov *et al.*, 1998, 2000a, b, 2001; Andrade *et al.*, 2003). They all employed flow cytometry

to determine bacterial abundance and reported numbers ranging from 3.7×10^4 to 5.5×10^8 cells ml^{-1} . This range resembled that obtained in the present study. Reduced nutrient concentrations in surface waters account for the low bacterial abundance and the dominance of LNA cells (80%) there (Andrade *et al.*, 2003). Alves *et al.* (2014) studied the same sampling site of the South-west Atlantic Ocean as that of the present investigation and found that temperature, dissolved organic carbon (DOC) and depth strongly influence microbial abundance and diversity. They also discovered that microbial genes and metabolic pathways are stratified in the South-west Atlantic Ocean water column.

Microbial plankton biomass was dominated by heterotrophic bacteria (up to 85% in TW, and almost 100% in the lower water masses), with a small contribution (7–18%) of *Prochlorococcus* and *Synechococcus* in the surface waters only. Total (both auto- and heterotrophic) bacterial numbers decreased with depth, even though the cell size of individuals increased with depth (Buitenhuis *et al.*, 2012). The increase of the availability of trophic resources (carbon, nitrogen and phosphorus) with depth was followed by the development of bigger cells with higher metabolic rates, as we observed with the increase of relative numbers of HNA bacteria. At 500–800 m, there were similar proportions of HNA and LNA bacteria, but at depths >1000 m, the relative quantity of HNA bacteria increased. The HNA play important roles in microbial metabolism (Lebaron *et al.*, 2001; Vila-Costa *et al.*, 2012). In addition, most deep-sea bacteria have expressive amounts of rRNA and contribute to bathypelagic metabolism (Karner *et al.*, 2001; Herndl *et al.*, 2005). These findings are consistent with the elevated metabolic rates detected in bathypelagic microbes sampled near the seafloor (Nagata *et al.*, 2000). Taken together, these facts help explain the dominance of HNA bacteria in the deep sea.

The dominance of HNA bacterial cells at deep waters can also be attributed to a significant decrease of 2–3 orders of magnitude in the abundance of flagellates, ciliates and mesozooplankton in comparison with the surface (Tanaka & Rassoulzadegan, 2002; Koppelman *et al.*, 2005). The lack of predation pressure could favour the establishment of bigger bacterial cells, usually the most preyed size fraction (Jürgens & Güde, 1994). In the bathypelagic ocean conditions for sustaining the dominance of HNA bacteria were found: abundant supply of inorganic nutrients and detritic carbon, and small predation pressure. It is very probable that the bathypelagic bacterioplankton is controlled by bottom-up mechanisms. We did not access other microbial components as ciliates and flagellates, groups that are known being at very low abundances at 2000 m (Tanaka & Rassoulzadegan, 2002; Koppelman *et al.*, 2005).

Diurnal differences in mesozooplankton and larval fish abundance have been reported in several oceanic studies (Olivar & Sabatés, 1997; Thurman & Burton, 2001; Munk *et al.*, 2015). Relative to deeper waters, most organisms, especially phytoplankton, occur on the surface layer, probably because of the comparatively rich food supplies there (Fernández-Álamo & Färber-Lorda, 2006). In the study area, although the highest densities were observed at night-time near the surface, they did not differ significantly from those observed during the daytime. This fact can be attributed to differences in the number of samples collected from each water mass during the day and night periods. Diel variations in epipelagic (0–200 m) mesozooplankton were not detected, most

likely because of the distance between sampling depths (1 m vs 250 m). Sampling at discrete depths would improve our assessment of vertical migration, particularly for very small organisms. A study in the Irish Sea found that >70% of the zooplankton were distributed within the top 10 m on average, and were considered weak or non-migrating (Irigoién *et al.*, 2004). On the other hand, larger zooplankton tend to be more effective at vertical migration than smaller ones (Hayes *et al.*, 2001) and cover greater distances in the water column.

We found no significant differences between day and night samplings in terms of the dominant copepod groups. Nevertheless, large migratory copepods like *Calanoides carinatus* and *Rhincalanus cornutus* were observed in the deep waters (SACW, AAIW and UCDW). Copepod dominance is reported in most mesozooplankton studies and demonstrates the importance of this group in transferring energy between different trophic levels (Turner, 2004; Escribano *et al.*, 2009; de Lira *et al.*, 2014; Munk *et al.*, 2015).

The copepod species identified in this study occur throughout all oceanic epipelagic, mesopelagic and bathypelagic zones (Cavalcanti & Larrazábal, 2004; Razouls *et al.*, 2005–2017; Bonecker, 2006; Lopes *et al.*, 2006; Dias *et al.*, 2010; Brugnano *et al.*, 2012; Bonecker *et al.*, 2014b). *Nullosetigera impar*, newly discovered in Brazilian waters, is mesopelagic to bathypelagic and was, until recently, only detected in the waters of the Central Atlantic, North-eastern Pacific and Indian Oceans (Devevey & Brooks, 1977; Razouls *et al.*, 2005–2017). The novel findings of the present study may be explained by the relative lack of prior investigation into the deep waters of the South Atlantic. These findings also underscore the fact that zooplankton richness in our area is underestimated (Bonecker *et al.*, 2014b).

The relative contributions of large copepods (*Candacia* spp. and *Pleuromamma* spp.) increased in night-time shallow waters and in deep waters. This pattern was also observed in the waters of the French Atlantic coast (Maycas *et al.*, 1999) and the Western Mediterranean (Brugnano *et al.*, 2012) and reflects the feeding habits of these organisms. Small omnivorous/herbivorous copepods tend to concentrate in the surface layer whereas larger detritivores/carnivores are usually found in deep waters. Large copepods feed and defecate throughout the water column, thereby regulating upward and downward carbon and nitrogen transfers (Maycas *et al.*, 1999). Maycas *et al.* (1999) noted that on the French Atlantic coast, larger copepods occurred at lower depths than the smaller ones. The authors proposed the small copepods migrated little or not at all. Ohman & Romagnan (2016) found that small non-migratory copepods remain in shallow waters both night and day whereas larger non-migratory organisms stay deeper in subsurface waters.

We found large densities of organisms belonging to the genus *Oithona*. These organisms were responsible for the formation of assemblages in several water masses. Because of their small size, organisms of the genus *Oithona* were probably underestimated in the present study because of the mesh size used. In a study carried out in the South Atlantic comparing nets of 60, 100 and 330 µm mesh sizes, the 100 µm net had the highest efficiency; thus, this mesh is more suitable for sampling zooplankton, and is more efficient for collecting small organisms, including representatives of the genus *Oithona* (Makabe *et al.*, 2012). In comparison, other studies have shown the importance of organisms from the

genus *Oithona* when using mesh sizes larger than 100 µm. In the waters off north-east Brazil, the species of this genus were qualitatively representative in samples collected with the net of 300 µm mesh size (Cavalcanti *et al.*, 2008). In a study developed in the Arctic region, where organisms collected with a net of 180 µm mesh size were identified, species of the genus *Oithona* dominated zooplankton assemblages (Gluchowska *et al.*, 2017). Another study carried out in the Campos Basin showed that *Oithona* species collected with a net of 200 µm mesh size were frequent in several water masses, and were responsible for the formation of communities in some depths (Bonecker *et al.*, 2014b). The great abundance of *Oithona* in our samples and the results obtained in the published literature confirm that *Oithona* are representatives of microzooplankton (20–200 µm), in addition to being important components of mesozooplankton (>200 µm). The present study aimed to collect several zooplankton groups, especially the largest mesozooplankton fractions; for this reason, we used a net of 200 µm mesh size, which is more suitable for these organisms (Sameoto *et al.*, 2000).

In this study, the densities of Euphausiacea, Chaetognatha, Doliolida and Salpida were significantly higher in night-time TW than the other water masses. The most abundant species in these groups are well-known migrators, including the euphausiids *Euphausia americana*, *E. similis* and *Nematoscelis atlantica*, the chaetognath *Flaccisagitta enflata*, and the salpids *Salpa fusiformis* and *Thalia democratica* (Mauchline, 1980; Hirota *et al.*, 1984; Madin *et al.*, 1996; Resgalla *et al.*, 2004; Lie *et al.*, 2012; Nogueira *et al.*, 2015).

The densities of Branchiopoda and Appendicularia did not significantly differ among the water masses in both sampling periods. The epipelagic *Pseudevadne tergestina* was detected in the top 100 m on the French Atlantic coast in the daytime (Maycas *et al.*, 1999) and in the coastal subtropical area (Miyashita *et al.*, 2011). The appendicularians *Oikopleura cornutogastra* and *Fritillaria formica* were found from the surface down to 2300 m in the Campos Basin, Brazil (Bonecker *et al.*, 2014b).

There was no significant difference between the day and night periods in terms of mollusc larva abundance. Nevertheless, the highest counts were obtained for the night-time TW and resemble those reported by Garland *et al.* (2002). The authors stated that molluscs are concentrated near the surface at night to take advantage of increased food source availability. The high mollusc densities observed in our study confirm the dispersal ability of these organisms from the coastal region to the oceanic region, which lowers their risk of extinction (Sahara *et al.*, 2015).

The mesopelagic fish *Lepidophanes guentheri* was among those that grouped the night-time TW samples. Many myctophids undergo daily vertical migration and enrich the carbon stocks in deep waters as they feed on the surface and defecate in the mesopelagic and bathypelagic zones (Angel, 2003; Conley & Hopkins, 2004; Castro *et al.*, 2010; Ariza *et al.*, 2015 and references within). *Lepidophanes guentheri* and *Lepidophanes gausi* migrate at night from the mesopelagic zone to the epipelagic zone, moving from depths of 700–950 and 425–850 m, respectively (Nafpaktitis *et al.*, 1977; Richards, 2006; Santos & Figueiredo, 2008).

In this study, it was found that *Cyclothone braueri* larvae were widely distributed throughout the water column. In a study of the Sargasso Sea, Sutton *et al.* (2010) reported that

the larvae of this species were the most abundant. Samples were collected from 0–1000 m (47.5%) and from 1000–5000 m (41.0%).

The indicator species *E. americana*, *E. similis*, *N. atlantica* and *Pterosagitta draco* are usually found in surface waters (Sameoto *et al.*, 1987; Pierrot-Bults & Nair, 1991). *Nannocalanus minor* is considered an indicator of the Brazil Current (Dias *et al.*, 2010) and is concentrated in the lower strata down to ~200 m (Björnberg, 1981; Cavalcanti & Larrázabal, 2004). The presence of *L. gaussi* and *L. guentheri* in night-time TW can be explained by the fact that both of them undergo nocturnal vertical migration as previously discussed in this article. Indicator species analysis (ISA) is an important tool in mesozooplankton ecology evaluation. Nevertheless, it must be used with caution when regarding strong migratory species because the data may reflect diel migratory behaviour of the organisms only during a period of the day, as in this study. The fact that certain species known to occur at great depths are used as night-time TW indicators is evidence of their daily migration.

In conclusion, only a portion of the mesozooplankton and larval fish communities undergo diel vertical migration. These include the euphausiids *E. americana*, *E. similis* and *N. atlantica*, and the larval fish *L. guentheri*. To the best of our knowledge, this study was the first attempt to describe the daily vertical distribution of mesozooplankton and larval fish communities in an oceanic region of the Brazilian coast. We furnished new data and insights on deep water vertical distribution and reported the occurrence of a new species in the heretofore poorly explored South-western Atlantic Ocean. Our results showed significant variability in mesozooplankton, larval fish and bacterioplankton abundance and distribution along an oceanic water column. Further studies are required to assess the factors driving diel vertical migration in this oceanic region. In this research, more frequent daily sampling at narrower depth ranges is required, and all water masses and plankton trophic levels should be considered in the process.

ACKNOWLEDGEMENTS

This study is part of the Habitats Project – Campos Basin Environmental Heterogeneity by CENPES/PETROBRAS. We thank the team at the Zooplankton and Ichthyoplankton Integrated Laboratory of the Federal University of Rio de Janeiro for sorting the samples. We also thank the following researchers for their help with identifications: Michele Arruda (Chaetognatha); Paulo R.F.C. Costa and Marta C.C. Quintas (Branchiopoda); Suzanna C. Vianna (Copepoda); Patrícia Alpino, Lohengrin Fernandes, Ralf Schawmborn, Cíntia Cardoso and Ana Cecilia R. Resende (decapod larvae); and Andrea S. Freire (Euphausiacea).

REFERENCES

Alves N. Jr., Meirelles P.M., de Oliveira S.E., Dutilh B., Silva G.G., Paranhos R., Cabral A.S., Rezende C., Lida T., de Moura R.L., Kruger R.H., Pereira R.C., Valle R., Sawabe T., Thompson C. and Thompson F. (2014) Microbial community diversity and physical-chemical features of the Southwestern Atlantic Ocean. *Archives of Microbiology* 197, 165–179.

- Andrade L., Gonzalez A.M., Araujo F.V. and Paranhos R. (2003) Flow cytometry assessment of bacterioplankton in tropical marine environments. *Journal of Microbiological Methods* 19, 89–94.
- Angel M.V. (2003) The pelagic environment of the open ocean. In Tyler P.A. (ed.) *Ecosystems of the deep oceans*, vol. 28. Amsterdam: Elsevier, pp. 39–79.
- Ariza A., Garijo J.C., Landeira J.M., Bordes F. and Hernández-León S. (2015) Migrant biomass and respiratory carbon flux by zooplankton and micronekton in the subtropical northeast Atlantic Ocean (Canary Islands). *Progress in Oceanography* 134, 330–342. doi: 10.1016/j.pocean.2015.03.003.
- Berasategui A.D., Menu Marque S., Gomez-Erache M., Ramirez F.C., Mianzan H.W. and Acha E.M. (2006) Copepod assemblages in a highly complex hydrographic region. *Estuarine, Coastal and Shelf Science* 66, 483–492.
- Björnberg T.K.S. (1981) Copepoda. In Boltovskoy D. (ed.) *Atlas del Zooplancton del Atlantico Sudoccidental y métodos de trabajo con el zooplancton marino*. Mar del Plata: INIDEP, pp. 587–679.
- Boebel O., Schmid C. and Zenk W. (1997) Flow and recirculation of Antarctic Intermediate Water across the Rio Grande Rise. *Journal of Geophysical Research* 102, 20967–20986.
- Boltovskoy D. (1999) *South Atlantic zooplankton*. Leiden: Backhuys Publishers, 1706 pp.
- Bonecker A.C.T. and Castro M.S. (eds) (2006) *Atlas de larvas de peixes da região central da Zona Econômica Exclusiva brasileira*. Museu Nacional Série Livros n. 19. Rio de Janeiro, Brasil. 216 pp.
- Bonecker A.C.T., Katsuragaewa M., Castro M.S., Gomes E.P., Namiki C.A.P. and Zani-Teixeira M. L. (2012) Larval fish of the Campos Basin, southeastern Brazil. *Check List* 8, 1280–1291.
- Bonecker A.C.T., Namiki C.A.P., Castro M.S. and Campos P.N. (2014a) *Catálogo dos estágios iniciais de desenvolvimento dos peixes da bacia de Campos* [online]. Curitiba: Sociedade Brasileira de Zoologia, 295 pp. Zoologia: guias e manuais de identificação series. ISBN 978-85-98203-10-2. Available from SciELO Books. <<http://books.scielo.org>>.
- Bonecker S.L.C. (ed.) (2006) *Atlas de zooplâncton da região central da Zona Econômica Exclusiva brasileira*. Museu Nacional Série livros n. 21. Rio de Janeiro, Brasil. 232 pp.
- Bonecker S.L.C., Araujo A.V., Carvalho P.F., Dias C.O., Loureiro Fernandes L.F., Migotto A.E. and Oliveira O.M.P. (2014b) Horizontal and vertical distribution of mesoplankton species richness and composition down to 2300 m in the southwest Atlantic Ocean. *Zoologia* 31, 445–462.
- Brugnano C., Granata A., Guglielmo L. and Zagami G. (2012) Spring diel vertical distribution of copepod abundances and diversity in the open Central Tyrrhenian Sea (Western Mediterranean). *Journal of Marine Systems* 105–108, 207–220.
- Buitenhuis E.T., Li W.K.W., Vaulot D., Lomas M.W., Landry M.R., Partensky F., Karl D.M., Ulloa O., Campbell L., Jacquet S., Lantoine F., Chavez F., Macias D., Gosselin M. and McManus G.B. (2012) Picoplankton biomass distribution in the global ocean. *Earth System Science Data* 4, 37–46.
- Campos E.J.D., Gonçalves J.E. and Ikeda Y. (1995) Water mass characteristics and geostrophic circulation in the South Brazil Bight: summer of 1991. *Journal of Geophysical Research* 100, 18537–18550.
- Campos E.J.D., Ikeda Y., Castro B.M., Gaeta S.A., Lorenzetti J.A. and Stevenson M.R. (1996) Experiment studies circulation in the Western South Atlantic. *Eos Transactions American Geophysical Union* 77, 253–259.
- Castro M.S., Richards W.J. and Bonecker A.C.T. (2010) Occurrence and distribution of larval lantern fish (Myctophidae) from the southwest Atlantic Ocean. *Zoologia* 27, 541–553.

- Cavalcanti E.A.H. and Larrazábal M.E.L.** (2004) Macrozooplâncton da Zona Econômica Exclusiva do Nordeste do Brasil (segunda expedição oceanográfica – REVIZEE/NE II) com ênfase em Copepoda (Crustacea). *Zoologia* 21, 467–475.
- Cavalcanti E.A.H., Neumann-Leitão S. and do Vieira D.A.N.** (2008) Mesozooplâncton do sistema estuarino de Barra das Jangadas, Pernambuco, Brasil. *Revista Brasileira de Zoologia* 25, 436–444.
- Cha S.S., McGowan M.F. and Richards W.J.** (1994) Vertical distribution of fish larvae off the Florida Keys, 26 May–5 June 1989. *Bulletin of Marine Science* 54, 828–842.
- Clarke K.R. and Warwick R.M.** (1994) *Change in marine communities: an approach to statistical analysis and interpretation*. Plymouth: N.E.R.C.
- Conley W.J. and Hopkins T.L.** (2004) Feeding ecology of lanternfish (Pisces: Myctophidae) larvae: prey preferences as a reflection of morphology. *Bulletin of Marine Science* 75, 361–379.
- Deevey G.B. and Brooks A.L.** (1977) Copepods of the Sargasso Sea off Bermuda: species composition, and vertical and seasonal distribution between the surface and 2000 m. *Bulletin of Marine Science* 22, 256–291.
- de Lira S.M.A., de Teixeira I.A., de Lima C.D.M., de Santos G.S., Neumann-Leitão S. and Schwamborn R.** (2014) Spatial and nyctemeral distribution of the zooneuston off Fernando de Noronha, Brazil. *Brazilian Journal of Oceanography* 62, 35–45.
- Dias C.O., Araujo A.V., Paranhos R. and Bonecker S.L.C.** (2010) Vertical copepod assemblages (0–2300 m) off Southern Brazil. *Zoological Studies* 49, 230–242.
- Dias C.O., Araujo A.V., Vianna S.C., Loureiro Fernandes L.F., Paranhos R., Suzuki M.S. and Bonecker S.L.C.** (2015) Spatial and temporal changes in biomass, production and assemblage structure of mesozooplanktonic copepods in the tropical south-west Atlantic Ocean. *Journal of the Marine Biological Association of the United Kingdom* 95, 483–496.
- Dufrêne M. and Legendre P.** (1997) Species assemblages and indicator species: the need for flexible asymmetrical approach. *Ecological Monographs* 67, 345–366.
- Escribano E., Hidalgo P. and Krautz C.** (2009) Zooplankton associated with the oxygen minimum zone system in the northern upwelling region of Chile during March 2000. *Deep-Sea Research Part II* 56, 1083–1094.
- Fernández-Álamo M.A. and Färber-Lorda J.** (2006) Zooplankton and the oceanography of the eastern tropical Pacific: a review. *Progress in Oceanography* 69, 318–359.
- Frost B. and Bollens S.** (1992) Variability of diel vertical migration in the marine planktonic copepod *Pseudocalanus newmani* in relation to its predators. *Canadian Journal of Fisheries and Aquatic Sciences* 49, 1137–1141.
- Gabioux M.** (2008) *Estudo numérico dos meandros e vórtices da Corrente do Brasil entre 22°S e 30°S*. PhD thesis. Rio de Janeiro: Universidade Federal do Rio de Janeiro, COPPE/UFRJ, 158 unpublished.
- Garland E.D., Zimmer C.A. and Lentz S.J.** (2002) Larval distributions in inner-shelf waters: the roles of wind-driven cross-shelf currents and diel vertical migrations. *Limnology and Oceanography* 47, 803–817.
- Gasol J.M. and del Giorgio P.A.** (2000) Using flow cytometry for counting natural planktonic bacteria and understanding the structure of planktonic bacterial communities. *Scientia Marina* 64, 197–224.
- Gluchowska M., Trudnowska E., Goszczko I., Kubiszyn A.M., Blachowiak-Samolyk K., Walczowski W. and Kwasniewski S.** (2017) Variations in the structural and functional diversity of zooplankton over vertical and horizontal environmental gradients en route to the Arctic Ocean through the Fram Strait. *PLoS ONE* 12, e0171715.
- Gonzalez-Silvera A., Santamaria-del-Angel E., Garcia V.M.T., Garcia C.A.E., Millán-Núñez R. and Muller-Karger F.** (2004) Biogeographical regions of the tropical and subtropical Atlantic Ocean off South America: classification based on pigment (CZCS) and chlorophyll-a (SeaWiFS) variability. *Continental Shelf Research* 24, 983–1000.
- Grasshoff K., Ehrhardt M. and Kremling K.** (1999) *Methods of seawater analysis*. Weinheim: Wiley-Verlag Chemie, 217 pp.
- Harris B.P., Young J.W., Reville A.T. and Taylor M.D.** (2014) Understanding diel-vertical feeding migrations in zooplankton using bulk carbon and nitrogen stable isotopes. *Journal of Plankton Research* 36, 1159–1163.
- Hays G.** (2003) A review of the adaptive significance and ecosystem consequences of zooplankton diel vertical migrations. *Hydrobiologia* 503, 163–170.
- Hays G.C., Harris R.P. and Head R.N.** (2001) Diel changes in the near-surface biomass of zooplankton and the carbon content of vertical migrants. *Deep-Sea Research II* 48, 1063–1068.
- Herndl G.J., Reinthaler T., Teira E., van Aken H., Veth C., Pernthaler A. and Pernthaler J.** (2005) Contribution of Archaea to total prokaryotic production in the deep Atlantic Ocean. *Applied and Environmental Microbiology* 71, 2303–2309.
- Hill A.E.** (1998) Horizontal zooplankton dispersal by diel vertical migration in S_2 tidal currents on the northwest European continental shelf. *Continental Shelf Research* 14, 491–506.
- Hirota Y., Nemoto T. and Marumo R.** (1984) Vertical distribution of larvae of *Euphausia nana* and *E. similis* (Crustacea: Euphausiacea) in Sagami Bay and Suruga Bay, central Japan. *Marine Biology* 81, 131–137.
- Irigoien X., Conway D.V.P. and Harris R.P.** (2004) Flexible diel vertical migration behaviour of zooplankton in the Irish Sea. *Marine Ecology Progress Series* 267, 85–97.
- Isla A., Scharek R. and Latasa M.** (2015) Zooplankton diel vertical migration and contribution to deep active carbon flux in the NW Mediterranean. *Journal of Marine Systems* 143, 86–97.
- Jackson D.A.** (1993) Stopping rules in principal components analysis – a comparison of heuristic and statistical approaches. *Ecology* 74, 2204–2214.
- Jung-Hoon K., Minho S., Youn K. and Woong-Seo K.** (2013) Diel vertical migration of the copepod *Calanus sinicus* before and during formation of the Yellow Sea Cold Bottom Water in the Yellow Sea. *Acta Oceanologica Sinica* 32, 99–106.
- Jürgens K. and Güde H.** (1994) The potential importance of grazing-resistant bacteria in planktonic systems: a review. *Marine Ecology Progress Series* 112, 169–188.
- Karner M.B., DeLong E.F. and Karl D.M.** (2001) Archaeal dominance in the mesopelagic zone of the Pacific Ocean. *Nature* 409, 507–510.
- Koppelman R., Timm H.T. and Weikert H.** (2005) Bacterial and zooplankton distribution in deep waters of the Arabian Sea. *Deep-Sea Research Part I* 52, 2182–2192.
- Krause M. and Radach G.** (1989) On the relations of vertical distribution, diurnal migration and nutritional state of herbivorous zooplankton in the Northern North Sea during FLEX 1976. *International Review of Hydrobiology* 74, 371–417.
- Lebaron P., Servais P., Agogue H., Courties C. and Joux F.** (2001) Does the high nucleic acid content of individual bacterial cells allow us to discriminate between active cells and inactive cells in aquatic systems? *Applied and Environmental Microbiology* 67, 1775–1782.

- Lie A.A.Y., Tse P. and Wong C.K. (2012) Diel vertical migration and feeding of three species of chaetognaths (*Flaccisagitta enflata*, *Aidanosagitta delicata* and *Aidanosagitta neglecta*) in two shallow, subtropical bays in Hong Kong. *Journal of Plankton Research* 34, 670–684.
- Lopes R.M., Katsuragawa M., Dias J.F., Montú M.A., Muelbert J.H., Gorri C. and Brandini F.P. (2006) Zooplankton and ichthyoplankton distribution on the southern Brazilian shelf: an overview. *Scientia Marina* 70, 189–202.
- de Macedo-Soares L.C.P., Garcia C.A.E., Freire A.S. and Muelbert J.H. (2014) Large-scale ichthyoplankton and water mass distribution along the South Brazil shelf. *PLoS ONE* 9: e91241.
- Madin L.P., Kremer P. and Hacker S. (1996) Distribution and vertical migration of salps (Tunicata, Thaliacea) near Bermuda. *Journal of Plankton Research* 18, 747–755.
- Makabe R., Tanimura A. and Fukuchi M. (2012) Comparison of mesh size effects on mesozooplankton collection efficiency in the Southern Ocean. *Journal of Plankton Research* 34, 432–436.
- Mauchline J. (1980) The biology of mysids and euphausiids. *Advances in Marine Biology* 18, 1–681.
- Maycas E.R., Bourdillon A.H., Macquart-Moulin C., Passelaigue F. and Patriti G. (1999) Diel variations of the bathymetric distribution of zooplankton groups and biomass in Cap-Ferret Canyon, France. *Deep-Sea Research Part II* 46, 2081–2099.
- Mayzaud P. and Pakhomov E.A. (2014) The role of zooplankton communities in carbon recycling in the ocean: the case of the Southern Ocean. *Journal of Plankton Research* 36, 1543–1556.
- McCune B. and Mefford M.J. (1999) PC-ORD. *Multivariate analysis of ecological data*. Version 4.0. Gleneden Beach, OR: MjM Software.
- McEwen G.F., Johnson M.W. and Folsom T.R. (1954) A statistical analysis of the performance of the Folsom plankton sample splitter, based upon test observations. *Theoretical and Applied Climatology* 7, 502–527.
- McGeoch M.A., Rensburg B.J.V. and Botes A. (2002) The verification and application of bioindicators: a case study of dung beetles in a savanna ecosystem. *Journal of Applied Ecology* 39, 661–672.
- Mémery L., Arhan M., Alvarez-Salgado X.A., Messias M.J., Mercier H., Castro C.G. and Rios A.F. (2000) The water masses along the western boundary of the south and equatorial Atlantic. *Progress in Oceanography* 47, 69–98.
- Miyashita L.K., Gaeta S.A. and Lopes R.M. (2011) Life cycle and reproductive traits of marine podonids (Cladocera, Onychopoda) in a coastal subtropical area. *Journal of Plankton Research* 33, 779–792.
- Müller T.J., Ikeda Y., Zangenberg N. and Nonato L.V. (1998) Direct measurements of Western Boundary Currents off Brazil between 20°S and 28°S. *Journal of Geophysical Research* 103, 5429–5437.
- Munk P., Nielsen T.G. and Hansen B.W. (2015) Horizontal and vertical dynamics of zooplankton and larval fish communities during mid-summer in Disko Bay, West Greenland. *Journal of Plankton Research* 37, 554–570.
- Nafpaktitis B.G., Backus R.H., Craddock J.E., Haedrich R.L., Robinson B.H. and Karnella C. (1977) Family Myctophidae. Part 7 of the Monograph Series. In Gibbs R.H. Jr. (ed.) *Fishes of the Western North Atlantic*. New Haven, CT: Yale University, pp. 13–299.
- Nagata T., Fukuda H., Fukuda R. and Koike I. (2000) Bacterioplankton distribution and production in deep Pacific waters: large-scale geographic variations and possible coupling with sinking particles fluxes. *Limnology and Oceanography* 45, 426–435.
- Niencheski L.F., Baumgarten M.G.Z., Roso R.H. and Bastos L.A.P. (1999) *Oceanografia química - levantamento bibliográfico e identificação do estado atual do conhecimento*. Rio de Janeiro: MMA, CIRM, FEMAR.
- Nogueira M. Jr., Brandini F.P. and Codina J.C.U. (2015) Diel vertical dynamics of gelatinous zooplankton (Cnidaria, Ctenophora and Thaliacea) in a subtropical stratified ecosystem (South Brazilian Bight). *PLoS ONE* 10: e0144161.
- Ohman M.D. and Romagnan J.B. (2016) Nonlinear effects of body size and optical attenuation on diel vertical migration by zooplankton. *Limnology and Oceanography* 61, 765–770.
- Olivar M.P. and Sabatés A. (1997) Vertical distribution of fish larvae in the north-west Mediterranean Sea in spring. *Marine Biology* 129, 289–300.
- Palomares-García R.J., Gómez-Gutiérrez J. and Robinson C.J. (2013) Winter and summer vertical distribution of epipelagic copepods in the Gulf of California. *Journal of Plankton Research* 35, 1009–1026.
- Pearre S. Jr. (2003) Eat and run? The hunger/satiation hypothesis in vertical migration: history, evidence and consequences. *Biological Reviews* 78, 1–79.
- Peterson R.G. and Stramma L. (1991) Upper-level circulation in the South Atlantic Ocean. *Progress in Oceanography* 26, 1–73.
- Pierrot-Bults A.C. and Nair V.R. (1991) Distribution patterns in chaetognaths. In Bone Q., Kapp H. and Pierrot-Bults A.C. (eds) *The biology of Chaetognaths*. Oxford: Oxford University Press, pp. 86–117.
- Razouls C., De Bovée F., Kouwenberg J. and Desreumaux N. (2005–2017) Diversity and geographic distribution of marine planktonic copepods. <http://copepodes.obs-banyuls.fr/en> (accessed 7 June 2016).
- R Development Core Team (2010) *R: A language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing. ISBN 3-900051-07-0. <http://www.r-project.org> (accessed 7 June 2016).
- Resgalla C. Jr., Carvalho J.L., Pereira F.J., Rörig L.R., Rodrigues-Ribeiro M., Tamanaha M.S. and Proença L.A.O. (2004) Migração vertical e taxas fisiológicas de *Thalia democratica* (Salpidae: Thaliacea) na Reserva Marinha do Arvoredo, Santa Catarina. *Notas técnicas Facimar* 8, 45–54.
- Rezende C.E., Andrade L., Suzuki M.S., Faro B.C.M.T., Gonzales A.S.M. and Paranhos R. (2007) Hidroquímica. In Valentin J.L. (ed.) *Características hidrobiológicas da região central da Zona Exclusiva brasileira (Salvador, BA ao Cabo de São Tomé, RJ)*. Brasília: Série Documentos REVIZEE-SCORE CENTRAL, pp. 31–60.
- Richards W.J. (2006) *Early stages of Atlantic fishes: an identification guide for the Western North Atlantic. Volume I. and Volume II*. Boca Raton, FL: CRC Press, 2640 pp.
- Rodrigues S.V., Marinho M.M., Jonck C.C.C., Gonçalves E.S., Brant V.F., Paranhos R., Curbelo M.P. and Falcão A.P. (2014) Phytoplankton community structures in shelf and oceanic waters off southeast Brazil (20°–25°S), as determined by pigment signatures. *Deep-Sea Research Part I* 88, 47–62.
- Rodriguez J.M., Cabrero A., Gago J., Guevara-Fletcher C., Herrero M., Hernandez de Rojas A., Garcia A., Laiz-Carrión R., Vergara A.R., Alvarez P., Piñero C. and Saborido-Rey F. (2015) Vertical distribution and migration of fish larvae in the NW Iberian upwelling system during the winter mixing period: implications for cross-shelf distribution. *Fisheries Oceanography* 24, 274–290.
- Roe H.S.J. (1974) Observations on the diurnal vertical migrations of an oceanic animal community. *Marine Biology* 28, 99–113.
- Sahara R., Fukaya K., Okuda T., Hori M., Yamamoto T., Nakaoka M. and Noda T. (2015) Larval dispersal dampens population fluctuation and shapes the interspecific spatial distribution patterns of rocky intertidal gastropods. *Ecography* 38, 1–9.

- Sameoto D., Guglielmo L. and Lewis M.K.** (1987) Day/night vertical distribution of euphausiids in the eastern tropical Pacific. *Marine Biology* 96, 235–245.
- Sameoto D., Wiebe P., Runge J., Postel L., Dunn J., Miller C. and Coombs S.H.** (2000) Collecting zooplankton. In Harris R., Wiebe P.H., Lenz J., Skjoldal H.R. and Huntley M. (eds) *ICES zooplankton methodology manual*. pp. 55–81.
- Santos A.P. and Figueiredo J.L.** (2008) *Guia de identificação dos peixes da família myctophidae do brasil*. São Paulo: Editora da Universidade de São Paulo, 168 pp.
- Silveira I.C.A.** (2007) O Sistema Corrente do Brasil na Bacia de Campos, RJ. PhD thesis. São Paulo: Universidade de São Paulo, USP, 181 pp. unpublished.
- Siokou I., Zervoudaki S. and Christou E.D.** (2013) Mesozooplankton community distribution down to 1000 m along a gradient of oligotrophy in the Eastern Mediterranean Sea (Aegean Sea). *Journal of Plankton Research* 35, 1313–1330.
- Sogawa S., Sugisaki H., Saito H., Okazaki Y., Ono T., Shimode S. and Kikuchi T.** (2016) Seasonal and regional change in vertical distribution and diel vertical migration of four euphausiid species (*Euphausia pacifica*, *Thysanoessa inspinata*, *T. longipes*, and *Tessarabrachion oculatum*) in the northwestern Pacific. *Deep-Sea Research Part I* 109, 1–9.
- Steinberg D.K. and Landry M.R.** (2017) Zooplankton and the ocean carbon cycle. *Annual Review of Marine Science* 9, 413–444.
- Sutton T.T., Wiebe P.H., Madin L. and Bucklin A.** (2010) Diversity and community structure of pelagic fishes to 5000 m depth in the Sargasso Sea. *Deep-Sea Research Part II* 57, 2220–2233.
- Suzuki M.S., Rezende C.E., Paranhos R. and Falcão A.P.** (2015) Spatial distribution (vertical and horizontal) and partitioning of dissolved and particulate nutrients (C, N and P) in the Campos Basin, Southern Brazil. *Estuarine, Coastal and Shelf Science* 166, 4–12.
- Tanaka T. and Rassoulzadegan F.** (2002) Full-depth profile (0–2000 m) of bacteria, heterotrophic nanoflagellates and ciliates in the NW Mediterranean Sea: vertical partitioning of microbial trophic structures. *Deep-Sea Research Part II* 49, 2093–2107.
- Thurman H.V. and Burton E.A.** (2001) *Introductory oceanography*. Englewood Cliffs, NJ: Prentice Hall.
- Tiberti R. and Iacobuzio R.** (2013) Does fish predation influence the vertical distribution of zooplankton in high transparency lakes? *Hydrobiologia* 709, 27–39.
- Turner J.T.** (2004) The importance of small planktonic copepods and their roles in pelagic marine food webs. *Zoological Studies* 43, 255–266.
- Van Rensburg B., McGeoch M.A., Chown S.L. and Jaarsveld A.S.** (1999) Conservation of heterogeneity among dung beetles in the Maputaland Centre of Endemism. *Biological Conservation* 88, 145–153.
- Vila-Costa M., Gasol J.M., Sharma S. and Moran M.A.** (2012) Community analysis of high- and low-nucleic acid-containing bacteria in NW Mediterranean coastal waters using 16S rDNA pyrosequencing. *Environmental Microbiology* 14, 1390–1402.
- Williamson C.E., Fischer J.M., Bollens S.M., Overholt E.P. and Breckenridge J.K.** (2011) Toward a more comprehensive theory of zooplankton diel vertical migration: integrating ultraviolet radiation and water transparency into the biotic paradigm. *Limnology and Oceanography* 56, 1603–1623.
- Yamaguchi A., Watanabe Y., Ishida H., Harimoto T., Furusawa K., Suzuki S., Ishizaka J., Ikeda T. and Takahashi M.M.** (2002) Community and trophic structures of pelagic copepods down to greater depths in the western subarctic Pacific (West-Cosmic). *Deep-Sea Research Part I* 49, 1007–1025.
- Zubkov M.V., Fuchs B.M., Burkill P.H. and Amann R.I.** (2001) Comparison of cellular and biomass specific activities of dominant bacterioplankton groups in stratified waters of the Celtic Sea. *Applied and Environmental Microbiology* 67, 5210–5218.
- Zubkov M.V., Sleight M.A. and Burkill P.H.** (2000a) Assaying picoplankton distribution by flow cytometry of underway samples collected along a meridional transect across the Atlantic Ocean. *Aquatic Microbial Ecology* 21, 13–20.
- Zubkov M.V., Sleight M.A., Burkill P.H. and Leakey R.J.G.** (2000b). Picoplankton community structure on the Atlantic Meridional Transect: a comparison between seasons. *Progress in Oceanography* 45, 369–386.
- and
- Zubkov M.V., Sleight M.A., Tarran G.A., Burkill P.H. and Leakey R.J.G.** (1998) Picoplanktonic community structure on a Atlantic transect from 50° N to 50° S. *Deep-Sea Research Part I* 45, 1339–1355.

Correspondence should be addressed to:

A. C. T. Bonecker
 Universidade Federal do Rio de Janeiro, Instituto de Biologia,
 Departamento de Zoologia, Av. Carlos Chagas Filho, 373 –
 Prédio do CCS, Bloco A, Sala Ao-084, Ilha do Fundão –
 21.941-902, Rio de Janeiro, RJ, Brasil
 email: ana@biologia.ufrj.br